MOD1 – DNA ENGINEERING

Bevin Engelward, Agi Stachowiak, David Weingeist

Writing Instructor: Neal Lerner

Oral Presentation Instructor: Atissa Banuazizi

Spring 2008


Day 6

The role of homologous recombination in rescuing cells from chemotherapeutics.


Culturing Mammalian Cells


Roles of Homologous Recombination:

Preventing Cancer
 & Affecting Treatment



BRCA2 Loads Rad51






Raw data from Grigorova et al., Cytogen. and Gen. Res. 104:333-340 (2004)


Homologous Recombination








Radiation & Chemotherapy

Replication Fork Animation By Tet Matsuguchi












RNA Polymerase




Transcription Factor










Expression is Off









Expression is On



Expression On



Add Doxycycline (= a tetracycline analog)



Expression On


tetO/P_{CMV}

Transcription

Rad51

Expression Off

Culturing Mammalian Cells







How do you grow mammalian cells?

What do cells need to grow in culture?

Correct Temperature

Correct pH Growth Factors (usually in

Salts

Correct Osmolality serum)

Amino Acids

Antibiotics

Glutamine (used for energy)

Lipids (usually in serum)

Vitamins

Glucose

Serum: Calf, Fetal, Horse, Bovine...

Typical media...

82 ml DMEM

15 ml Calf Serum

1.5 ml Glutamine

1.5 ml Penn/Strep

What is in DMEM?

Vitamins

Sa	lts
Sa	IlΟ

Amino Acids

	D 0422	VITAMINS	<u> </u>
	[1X]	Choline Bitartrate	_
COMPONENT	g/L	Choline Chloride	0.004
		Felic Acid	0.004
INORGANIC SALTS		myo-Inositol	0.0072
CaCl ₂ -2H ₂ O	0.265	Niacinamide	0.004
Fe(NO ₃) ₃ -9H ₂ O	0.0001	D-Pantothenic Acid+%Ca	0.004
MgSO ₄	0.09767	Pyridoxal·HCl	_
KCl	0.4	Pyridoxine*HCl	0.004
NaHCO ₃	3.7	Riboflavin	0.0004
NaCl	6.4	Thiamine-HCl	0.004
NaH ₂ PO ₄	0.109	OTHER	
Succinic Acid		D-Gincose	4.3
Sodium Succinate		HEPES	_
AMINO ACIDS		Phenol Red Na	0.0159
L-Arginine-HCl	0.084	Pyruvic Acid-Na	0.11
L-Cystine-2HCl	_		
L-Glutamine	_	ADD	
Glycine	0.03	Glucose	_
L-Histidine+HCl+H ₂ O	0.042	L-Ghtamine	0.584
L-Isoleucine	0.105		
L-Leucine	0.105	L-Cystine 2HCl	-
L-Lycine HCl	0.146	L-Leucine	_
L-Methionine		L-Lysine•HCl	-
L-Phenylalanine	0.066	L-Methionine	-
L-Serine	0.042	NaHCO ₃	_
L-Threquine	0.095	NaH ₃ PO ₄	_
L-Tryptophan	0.016	Phenol Red Na	-
L-Tyrosine (free base)	 	Pyruvic Acid•Na	_
L-Tyrosine-2Na-2H ₂ O	0.10379		
L-Valine	0.094	Grams of powder required to prepare 1 L	N/A

How do you maintain a neutral pH?

Blood pH is 7.4

The most important buffer in extracellular fluids is a mixture of carbon dioxide (CO_2) and bicarbonate anion (HCO_3^-) .

$$H_2O + CO_2 \leftrightarrow H_2CO_3 \leftrightarrow H^+ + HCO_3^-$$

$$NaOH + H_2CO_3 \leftrightarrow NaHCO_3 + H_2O$$

$$NaHCO_3 \leftrightarrow Na^+ + HCO_3^-$$


The pH is determined by the concentration of CO₂ and bicarbonate.

Mammalian Cell Incubators





Why is sterility important?



From Freshney's "Culture of Mammalian Cells"

Mammalian Cell Culture Hood








Hands-On..


What does it mean to "pass" your cultures?

Cell Density (cells per cm²)










Low Density High Density

Confluent

Confluent cells are often contact inhibited

Cell Density can be kept low through "Passaging"




Trypsin is used to release cells from the surface of the dish












Cell Culture Growth Phases



Things you need to know about your cells...

Lag phase

Log phase

Doubling time

Optimal Maximum Density


Optimal Minimum Density


Growth inhibition – arrest

Plating Efficiency

Contact Inhibition

ATCC Number: CCL-92 3T3 Swiss Albino Designation:









Low Density High Density


Confluent



You need to know what your cells should look like!



Sick Cells:



The role of homologous recombination in rescuing cells from chemotherapeutics.

Culturing Mammalian Cells