



Cancer Research

Definition of a FoxA1 Cistrome That Is Crucial for G₁ to S-Phase Cell-Cycle Transit in Castration-Resistant Prostate Cancer

Chunpeng Zhang, Liguang Wang, Dayong Wu, et al.

Cancer Res 2011;71:6738-6748. Published OnlineFirst September 7, 2011.

Updated Version

Access the most recent version of this article at:
doi:[10.1158/0008-5472.CAN-11-1882](https://doi.org/10.1158/0008-5472.CAN-11-1882)

Supplementary Material

Access the most recent supplemental material at:
<http://cancerres.aacrjournals.org/content/suppl/2011/09/07/0008-5472.CAN-11-1882.DC1.html>

Cited Articles

This article cites 46 articles, 18 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/71/21/6738.full.html#ref-list-1>

E-mail alerts

[Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions

To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions

To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.

Definition of a FoxA1 Cistrome That Is Crucial for G₁ to S-Phase Cell-Cycle Transit in Castration-Resistant Prostate Cancer

Chunpeng Zhang¹, Liguang Wang⁵, Dayong Wu¹, Hongyan Chen¹, Zhong Chen¹, Jennifer M. Thomas-Ahner², Debra L. Zynger³, Jérôme Eeckhoutte⁶, Jindan Yu⁷, Jun Luo⁸, Myles Brown⁹, Steven K. Clinton², Kenneth P. Nephew¹⁰, Tim H.-M. Huang⁴, Wei Li⁵, and Qianben Wang¹

Abstract

The enhancer pioneer transcription factor FoxA1 is a global mediator of steroid receptor (SR) action in hormone-dependent cancers. In castration-resistant prostate cancer (CRPC), FoxA1 acts as an androgen receptor cofactor to drive G₂ to M-phase cell-cycle transit. Here, we describe a mechanistically distinct SR-independent role for FoxA1 in driving G₁ to S-phase cell-cycle transit in CRPC. By comparing FoxA1 binding sites in prostate cancer cell genomes, we defined a codependent set of FoxA1-MYBL2 and FoxA1-CREB1 binding sites within the regulatory regions of the Cyclin E2 and E2F1 genes that are critical for CRPC growth. Binding at these sites upregulate the Cyclin E2 and Cyclin A2 genes in CRPC but not in earlier stage androgen-dependent prostate cancer, establishing a stage-specific role for this pathway in CRPC growth. Mechanistic investigations indicated that FoxA1, MYBL2, or CREB1 induction of histone H3 acetylation facilitated nucleosome disruption as the basis for codependent transcriptional activation and G₁ to S-phase cell-cycle transit. Our findings establish FoxA1 as a pivotal driver of the cell-cycle in CRPC which promotes G₁ to S-phase transit as well as G₂ to M-phase transit through two distinct mechanisms. *Cancer Res*; 71(21); 6738–48. ©2011 AACR.

Authors' Affiliations: ¹Department of Molecular and Cellular Biochemistry and the Comprehensive Cancer Center, ²Division of Medical Oncology, Department of Internal Medicine and the Comprehensive Cancer Center, and ³Department of Pathology, The Ohio State University College of Medicine; ⁴Human Cancer Genetics Program, Department of Molecular Virology, Immunology, and Medical Genetics, The Ohio State University, Columbus, Ohio; ⁵Division of Biostatistics, Dan L. Duncan Cancer Center and Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, Texas; ⁶INSERM UMR 1011-Bâtiment J&K, Université Lille-Nord de France, Faculté de Médecine de Lille-Pôle Recherche, Lille cedex, France; ⁷Department of Medicine and Robert H. Lurie Comprehensive Cancer Center, Northwestern University Feinberg School of Medicine, Chicago, Illinois; ⁸Department of Urology, The Johns Hopkins University School of Medicine, Baltimore, Maryland; ⁹Department of Medical Oncology, Dana-Farber Cancer Institute and Harvard Medical School, Boston, Massachusetts; and ¹⁰Medical Sciences Program, Department of Cellular and Integrative Physiology, Indiana University School of Medicine, Bloomington, Indiana

Note: Supplementary data for this article are available at Cancer Research Online (<http://cancerres.aacrjournals.org>)

C. Zhang, L. Wang, and D. Wu contributed equally to this work.

Corresponding Authors: Qianben Wang, Department of Molecular and Cellular Biochemistry and the Comprehensive Cancer Center, The Ohio State University College of Medicine, 888 Biomedical Research Tower, 460 West 12th Ave, Columbus, OH 43210. Phone: 614-247-1609; Fax: 614-688-4181; E-mail: qianben.wang@osumc.edu; or Wei Li, Division of Biostatistics, Dan L. Duncan Cancer Center and Department of Molecular and Cellular Biology, Baylor College of Medicine, One Baylor Plaza, Cullen Building, Suite 450A9, Houston, TX 77030. Phone: 713-798-7854; Fax: 713-798-2716; E-mail: WL1@bcm.edu

doi: 10.1158/0008-5472.CAN-11-1882

©2011 American Association for Cancer Research.

Introduction

The FoxA subfamily of winged helix/forkhead box (Fox) transcription factors, which consists of 3 members, FoxA1, FoxA2, and FoxA3, have been found to play important roles in multiple stages of development, metabolism, differentiation, and proliferation (1). FoxA proteins function as "pioneer factors" that engage chromatin before other transcription factors (2–5). Recent studies have further shown that FoxA1 functions as a pioneer factor for steroid hormone receptors (SR), including androgen receptor (AR) in prostate cancer cells and estrogen receptor (ER) in breast cancer cells, directing AR- and ER-regulated hormone (androgen and estrogen)-responsive genes (6–8). These studies further suggested that FoxA1 acts upstream of AR and ER to regulate their target genes in hormone-dependent prostate and breast cancers.

In both androgen-dependent prostate cancer (ADPC) and fatal castration-resistant prostate cancer (CRPC), AR expression and functionality have been well documented (9), but the receptor seems to play different roles in the two diseases. For example, in ADPC, AR functions primarily to promote G₁–S cell-cycle progression, by transcriptional and/or posttranscriptional regulation of CDKN1A, CCND1, and CDKN1B (10). However, in CRPC, the primary function of AR seems to be regulation of G₂–M transition. Thus, the receptor seems to be "reprogrammed" to direct transcriptional regulation of

G₂-M phase-specific genes, including *UBE2C* and *CDK1* (11), by mechanisms that are not well understood. Consistent with the critical role of FoxA1 in assisting SR binding, FoxA1 silencing in CRPC leads to decreased AR binding to enhancers of G₂-M phase genes and lower gene expression levels (11). Hence, in CRPC, FoxA1 seems to be mainly involved in AR-regulated G₂-M cell-cycle progression. However, whether this pioneer factor can also function independently in CRPC, in addition to collaborating with AR, has not been investigated.

In this study, we examined the role of FoxA1 in CRPC cell-cycle progression by silencing FoxA1 in unsynchronized CRPC cells. We found that in the absence of FoxA1, CRPC experienced a G₁-S block and, unexpectedly, not G₂-M arrest. Our comprehensive integrated analysis of gene expression and FoxA1 cistrome data further revealed direct upregulation of *CCNE2* by FoxA1 binding sites specific to CRPC, as well as indirect upregulation of *CCNA2* by E2F1. We further established a requirement for CREB1 and MYBL2 in CRPC-specific FoxA1 binding, through histone H3 acetylation-facilitated nucleosome disruption, resulting in upregulation of *CCNE2* and *CCNA2* expression and enhanced growth of CRPC. Collectively, these data indicate that FoxA1, together with CREB1 and MYBL2, drive G₁-S progression in CRPC, which is distinct from the classic role of FoxA1 as an AR cofactor.

Materials and Methods

Cell lines

The ADPC cell line LNCaP was purchased from the American Type Culture Collection (ATCC) and cultured in RPMI 1640 media (Invitrogen) supplemented with 10% FBS. The CRPC cell line abl was provided by Zoran Culig (Innsbruck Medical University, Austria). A second CRPC cell line, CWR22Rv1, was provided by Steven P. Balk (Harvard Medical School, Boston, MA). LNCaP was authenticated by ATCC, and abl and CWR22Rv1 were authenticated as described previously (12, 13). All 3 cell lines were passaged in our laboratory for less than 6 months after resuscitation.

RNA interference

siRNA transfections were done by Lipofectamine 2000 (Invitrogen). The sequences for siRNAs were listed in Supplementary Table S1.

Western blots

Western blot analyses were carried out as previously described (14). Antibodies used are listed in the Supplementary Table S2.

Cell proliferation assay

Cell proliferation was measured by the WST-1 kit (Roche), as previously described (15).

Fluorescence-activated cell-sorting analysis

Cells were collected, and DNA contents were analyzed by a FACS Calibur cell flow cytometer (Becton Dickinson Biosciences) as previously described (14, 15).

Real-time reverse transcriptase PCR

Real-time reverse transcriptase (RT)-PCR was carried out as previously described (14). Primers used are listed in Supplementary Table S1.

ChIP-on-chip assay, ChIP assay, and re-ChIP assay

The ChIP-on-chip experiments were carried out in biological triplicates as previously described (11). The raw data of FoxA1 ChIP-on-chip have been submitted to the Gene Expression Omnibus (GEO) repository under the accession number GSE26329. Chromatin immunoprecipitation (ChIP) and re-ChIP assays were carried out as previously described (16). Antibodies for ChIP and re-ChIP assays are listed in Supplementary Table S2.

Correlation of FoxA1 cistrome in LNCaP and abl cells with clinical ADPC and CRPC microarray data

A meta-analysis was carried out by using 3 clinical ADPC/CRPC gene expression microarray data sets (17–19) from Oncomine (20). The overexpressed and underexpressed genes in CRPC versus ADPC, and randomly selected genes were correlated with 3 FoxA1 binding groups. Details are available in the Supplementary Materials and Methods.

Formaldehyde-assisted isolation of regulatory elements

Formaldehyde-assisted isolation of regulatory elements (FAIRE) was carried out as previously described (21, 22). All primer sequences are listed in Supplementary Table S1.

Motif finding in abl-specific FoxA1 binding regions

MDmodule (23) was used to conduct a *de novo* search for motifs significantly enriched in abl-specific binding sites with LNCaP-specific binding regions as background. Motif length varied from 6 to 15 nucleotides. Significantly enriched motif was annotated using STAMP (24).

Results

FoxA1 promotes CRPC cell growth by driving G₁-S and G₂-M cell-cycle progression

Our previous findings that FoxA1 binding at the enhancer regions of several cell-cycle G₂-M phase genes (e.g., *UBE2C* and *CDK1*) is required for overexpression of these genes in CRPC (11) prompted us to further examine the functional role of FoxA1 in CRPC. We first tested the effect of FoxA1 silencing on cell proliferation in abl, a CRPC cell line that closely models clinical CRPC (11, 12, 25). siRNAs targeting FoxA1 (2 independent siRNAs) were transfected into abl. Both siRNAs dramatically reduced cellular FoxA1 levels (Fig. 1A) and significantly decreased androgen-independent cell proliferation of abl (Fig. 1A). The inhibitory effect of FoxA1 silencing on cell proliferation was also observed in CWR22Rv1 (Supplementary Fig. S1A), another CRPC cell model (26). We next examined the effect of FoxA1 silencing on CRPC cell-cycle progression by using a thymidine-nocodazole block to enrich abl cells at mitosis (15). Consistent with the role of FoxA1 in the upregulation of cell-cycle G₂-M phase genes in CRPC cells (11), FoxA1 silencing caused a




Figure 1. FoxA1 silencing decreases CRPC cell growth by blocking G₂-M and G₁-S cell-cycle progression. A, abl cell proliferation was measured after siRNA transfection by using the WST-1 assay. **, $P < 0.01$. B, abl cells were transfected with siRNAs and synchronized in mitosis by using a thymidine-nocodazole block, and cells were analyzed by flow cytometry. *, $P < 0.05$; **, $P < 0.01$. C, 72 hours after siRNA transfection, abl cells were analyzed by flow cytometry. *, $P < 0.05$; **, $P < 0.01$.

G₂-M accumulation in the treated abl cells (Fig. 1B). However, FoxA1 knockdown resulted in a G₁-S block in unsynchronized abl and CWR22Rv1 cells (Fig. 1C; Supplementary Fig. S1B), suggesting that FoxA1 promotes G₁-S and G₂-M progression in CRPC cells. As a major role of AR in CRPC is to upregulate G₂-M phase gene expression and promote the G₂-M transition (11), these data further indicate that the ability of FoxA1 to promote G₁-S progression is an AR-independent phenomenon. In agreement with our previous findings (16), silencing of FoxA1 had no effect on dihydrotestosterone (DHT)-stimulated LNCaP G₁-S progression and cell proliferation (Supplementary Fig. S1C and D). Taken together, these data suggest that FoxA1 is required for androgen-independent but not androgen-dependent growth of prostate cancer.

FoxA1 upregulates G₁ phase genes *CCNE2* and *CCNA2* to enhance CRPC cell growth

To identify FoxA1-regulated genes that contribute to FoxA1-mediated G₁-S transition in CRPC cells, abl cells were transfected with siFoxA1 and a control siRNA (siControl), cultured for 72 hours, and analyzed for expression of the G₁ phase genes *CCNE2*, *CCNA2*, *CCNE1*, *CDK2*, *CDK4*, *CDK6*, *CCND1*, *CCND2*, *CCND3*, *CDKN1A*, and *CDKN1B* and *RB*. LNCaP cells transfected with these siRNAs and treated with DHT (10 nmol/L) for 4 hours and 24 hours served as controls. Among the 11 genes examined by real-time RT-PCR, expres-

sion levels of *CCNE2*, *CCNA2*, and *CDK2* were higher in abl cells transfected with siControl than similarly transfected, untreated or DHT-treated LNCaP (Fig. 2A), although *CCNE2*, *CCNA2*, and *CDK2* expression levels were increased after DHT treatment of LNCaP cells (Fig. 2A; ref. 27). Silencing of FoxA1 markedly decreased *CCNE2* and *CCNA2* but not *CDK2* expression in abl cells, but not LNCaP cells (Fig. 2A). Importantly, reanalysis of gene expression profiles from 3 independent clinical studies (17–19) confirmed that expression of *CCNE2* and *CCNA2* was greater in cases of CRPC than in cases of ADPC (Fig. 2B). Western blot analysis showed that *CCNE2* and *CCNA2* protein levels were also greater and were FoxA1-dependent in abl compared with LNCaP cells, in the presence or absence of DHT (Fig. 2C). Consistent with previous reports showing that *CCNE2* and *CCNA2* binding and activation of *CDK2* lead to RB phosphorylation (28, 29), FoxA1-enhanced expression of *CCNE2* and *CCNA2* protein increased FoxA1-dependent expression of phosphorylated *CDK2* and phosphorylated RB in abl versus LNCaP (Fig. 2C). The inhibitory effect of FoxA1 depletion on mRNA and protein expression levels of *CCNE2*, but not *CCNA2*, was also observed in CWR22Rv1 cells (Supplementary Fig. S2A).

We next examined the functional role of *CCNE2* and *CCNA2* in ADPC and CRPC cell growth. In cell proliferation assays, *CCNE2* or *CCNA2* silencing markedly decreased abl and CWR22Rv1 growth (Fig. 2D; Supplementary Fig. S2B), suggesting that *CCNE2* and *CCNA2* play an essential role in CRPC cell




Figure 2. FoxA1 promotes CRPC growth through upregulation of CCNE2 and CCNA2. A, 72 hours after siRNA transfection into LNCaP in the absence (–) or presence (+) of DHT for 4 hours and 24 hours, and abl cells in the absence of DHT, total RNA was isolated and amplified by real-time RT-PCR. The mRNA expression data are summarized in heat maps. The color scale represents relative mRNA expression level. B, boxplots show that CCNE2 and CCNA2 are overexpressed in CRPC cases (red) compared with ADPC cases (black). Number of patients is indicated in brackets. C, Western blot analyses were carried out by the indicated antibodies. D, abl cells were transfected with siRNAs in the absence of DHT, and LNCaP cells were transfected with siRNAs in the absence (–) or presence (+) of 10 nmol/L DHT. The cell proliferation was measured by the WST-1 assay. **, $P < 0.01$. E, LNCaP cells stably expressing pcDNA3.1-CCNE2, pcDNA3.1-CCNA2, or the empty vector were cultured in hormone-depleted medium and treated with DHT from 10^{-12} to 10^{-7} mol/L. The cell proliferation was measured by the WST-1 assay. **, $P < 0.01$ as compared with day 0.

proliferation. To test whether differences in expression of CCNE2 and CCNA2 were responsible for androgen-independent and androgen-dependent phenotypic differences between abl and LNCaP cells, we generated LNCaP cell lines stably expressing CCNE2 or CCNA2. Consistent with a previous report that overexpression of G₁ cyclins was not sufficient to induce LNCaP growth in the absence of androgen (30), no effect of CCNE2 or CCNA2 overexpression on cell proliferation and G₁–S progression was observed for vehicle-treated LNCaP cells (Fig. 2E; Supplementary Fig. S2C). However, while DHT treatment (1 nmol/L or higher) increased growth of vector transfected LNCaP, a growth response in LNCaP cells overexpressing CCNE2 or CCNA2 was observed at a lower (0.1 nmol/L) concentration of DHT (Fig. 2E), suggesting that

CCNE2 and CCNA2 overexpression increases LNCaP androgen sensitivity.

A distinct FoxA1 CRPC cistrome regulates differentially expressed genes in clinical CRPC versus ADPC

To investigate the mechanism underlying FoxA1-regulated CCNE2 and CCNA2 gene expression, we mapped the FoxA1 cistromes in LNCaP and abl cells by combining ChIP with Affymetrix human whole genome tiling arrays (ChIP-on-chip). Using the MAT (model-based analysis of tiling-array) algorithm (31) with a P value cut-off of $1E-4$ or less, we identified 14,965 and 18,110 FoxA1 binding sites in LNCaP and abl cells, respectively. Overlapping analysis of FoxA1 binding in LNCaP and abl identified 14,248 common FoxA1 binding regions

between the 2 cell lines, 717 LNCaP-specific FoxA1 binding sites and 3,862 abl-specific FoxA1 binding sites (Fig. 3A; Supplementary Fig. S3A). The ChIP-on-chip results were confirmed by using direct ChIP for FoxA1 on a subset of common FoxA1 binding regions, LNCaP-specific FoxA1 binding regions, abl-specific FoxA1 binding regions, and negative regions (Supplementary Fig. S3B). To determine the functional significance of common LNCaP-specific and abl-specific FoxA1 binding sites, these 3 types of FoxA1 binding sites were correlated to gene expression profiles from 3 clinical studies on ADPC and CRPC cases (17–19). Interestingly, abl-specific FoxA1 binding sites were significantly enriched within 20 or 40 kb of the transcription start sites (TSS) of overexpressed (e.g., *CCNE2*; Fig. 2B) and underexpressed genes (but more notably overexpressed genes) in CRPC versus ADPC, but not randomly selected genes (Fig. 3B; Supplementary Fig. S3C and D). By contrast, no obvious enrichment of LNCaP-specific FoxA1 binding sites and common FoxA1 binding sites was observed near overexpressed, underexpressed, or randomly selected genes (Fig. 3B; Supplementary Fig. S3C and D). These results suggest that abl-specific FoxA1 binding sites may, in general, directly regulate differentially expressed genes in CRPC versus ADPC.

Specific FoxA1 binding sites directly upregulate *CCNE2* in CRPC cells

On the basis of strong correlation between global FoxA1 binding and differential gene expression profiles, we next investigated regulation of *CCNE2* by FoxA1 in CRPC-abl and ADPC-LNCaP cells. ChIP-on-chip analysis identified an

abl-specific FoxA1 binding site at the *CCNE2* promoter region, 3 abl-specific FoxA1 binding sites located -24.0 , -9.8 , and $+23.5$ kb away from the TSS of *CCNE2*, and a common FoxA1 binding site (defined as MAT-score ≥ 3.72 in both cell lines; see the Supplementary Materials and Methods) 14.3 kb downstream of TSS of *CCNE2* (Fig. 4A). The common site displayed stronger FoxA1 binding in abl (MAT-score = 9.87) compared with LNCaP (MAT-score = 4.37). No overlap was seen between these abl-specific FoxA1 binding sites and AR binding regions in abl cells (ref. 11; data not shown). Direct ChIP analysis showed higher FoxA1 occupancy at the *CCNE2* promoter region and the 4 putative *CCNE2* enhancer regions in abl versus LNCaP cells (Fig. 4B). To further characterize the *CCNE2* promoter and the putative *CCNE2* enhancer regions, ChIP assays were carried out in LNCaP and abl cells, using antibodies against an enhancer histone mark H3K4 monomethylation (H3K4me1; ref. 32), phosphorylated RNA polymerase II at serine 5 (p-Pol II), and transcription coactivators CREB binding protein (CBP) and Mediator 1 (MED1). Enrichment of H3K4me1 was observed at the 4 putative *CCNE2* enhancer regions compared with the *CCNE2* promoter region in abl, and the level of H3K4me1 was higher at putative *CCNE2* enhancers 1, 2, and 4 in abl compared with LNCaP (Fig. 4C), indicating that these 4 distal FoxA1 binding regions function as enhancers in abl cells. Greater p-Pol II level and increased CBP and MED1 binding at the *CCNE2* enhancer and promoter regions in abl versus LNCaP cells (Fig. 4C) further supported the hypothesis that these FoxA1 binding sites may play more important transcriptional regulatory roles in abl than in LNCaP cells. In addition, H3K4me1 levels were higher, and




Figure 3. Correlation of CRPC cell-specific FoxA1 binding sites with clinical CRPC/ADPC gene expression data sets. **A**, overlap analysis showing the number of FoxA1 binding sites specific to LNCaP or abl or shared between the 2 cell lines ($P \leq 1E-4$). **B**, correlation between cell type-specific FoxA1 binding sites and differentially expressed genes in clinical CRPC versus ADPC. FoxA1 binding sites are divided into 3 groups: abl-specific (black), LNCaP-specific (white), and common binding sites between the 2 cell lines (gray). Three gene lists are presented as separated stacked bars along x-axis. P values on each bar show the significance level of association between FoxA1 binding and genes having FoxA1 binding sites with 20 kb of the TSS.




Figure 4. FoxA1 directly regulates *CCNE2* in CRPC cells. A, FoxA1 binding sites in LNCaP and abl cells relative to the *CCNE2* gene are shown, using the UCSC genome browser format. B and C, ChIP assays were carried out in LNCaP and abl cells in the absence of androgen, using an anti-FoxA1 antibody (B), or antibodies against H3K4me1, p-Pol II, CBP, or MED1 (C). *, $P < 0.05$; **, $P < 0.01$. D, LNCaP and abl cells were transfected with siRNAs in the absence of androgen. FAIRE-qPCR was carried out 72 hours posttransfection. *, $P < 0.05$; **, $P < 0.01$.

greater recruitment of FoxA1, p-Pol II, CBP, and MED1 at UBE2C and/or CDK1 enhancers was observed in abl versus LNCaP cells (Supplementary Fig. S4A), in agreement with our previous findings (11).

As transcriptionally active *cis*-regulatory elements always reside within nucleosome-depleted regions (NDR; ref. 33) and the FAIRE technique has been used to successfully identify NDRs (21, 22), we used FAIRE to analyze the local chromatin structure of the FoxA1 binding regions. LNCaP and abl cells were transfected with siControl or siFoxA1 followed by FAIRE. In siControl transfected cells, a higher FAIRE signal was observed at the *CCNE2* enhancer and promoter regions in abl cells than in LNCaP cells (Fig. 4D), and importantly, FoxA1 silencing decreased FAIRE enrichment at the *CCNE2* enhancer and promoter regions only in abl cells (Fig. 4D). Taken together, these data strongly indicate that increased FoxA1 binding induces nucleosome depletion at the *CCNE2* enhancer and promoter regions. We obtained essentially similar results at the UBE2C and CDK1 enhancers (Supplementary Fig. S4B).

CRPC-specific FoxA1 binding sites upregulate *CCNA2* via *E2F1*

Although, in general, FoxA1 binding was significantly enriched near overexpressed genes in CRPC compared with

ADPC (Fig. 3B), we were unable to identify, using the ChIP-on-chip technique, abl-specific FoxA1 binding sites near *CCNA2* (Fig. 5A), indicative of an indirect mechanism of FoxA1-mediated *CCNA2* upregulation in abl cells (Fig. 2A). As previous studies in other systems showed direct regulation of *CCNA2* by E2F1 transcription factor binding to the *CCNA2* promoter region (34, 35) and our FoxA1 ChIP-on-chip and direct FoxA1 ChIP analyses identified and confirmed 2 putative E2F1 enhancers reside 15.8 and 33.4 kb downstream of the TSS of *E2F1* (Fig. 5A and B), we hypothesized that FoxA1 may regulate *CCNA2* expression through a direct upregulation of *E2F1*. As expected, increased levels of H3K4me1 and p-Pol II, higher occupancy of CBP and MED1, and higher FAIRE enrichment at E2F1 enhancers, were observed in abl compared with LNCaP cells (Fig. 5C and D), resulting in FoxA1-dependent increased mRNA and protein levels of E2F1 in abl versus LNCaP cells in the absence of androgen (Fig. 5E). Similar to *CCNE2* regulatory regions, no AR binding was observed at these 2 E2F1 enhancers in abl cells (data not shown). We next performed ChIP to examine whether E2F1 directly regulates *CCNA2* in abl cells, and expression of *CCNA2* mRNA following E2F1 silencing was also examined in LNCaP and abl cells. As shown in Figure 5F and G, E2F1 binding at the *CCNA2* promoter was increased, and E2F1-dependent, *CCNA2*




Figure 5. FoxA1 regulates *CCNA2* via direct regulation of *E2F1*. **A**, FoxA1 binding sites in LNCaP and abl cells relative to the *CCNA2* and *E2F1* genes are shown, using the UCSC genome browser format. **B** and **C**, ChIP assays were carried out in LNCaP and abl cells in the absence of androgen, using an anti-FoxA1 antibody (**B**), or antibodies against p-Pol II, CBP, H3K4me1, or MED1 (**C**). *, $P < 0.05$; **, $P < 0.01$. **D**, LNCaP and abl cells were transfected with siRNAs in the absence of androgen. FAIRE-qPCR was carried out 72 hours posttransfection. *, $P < 0.05$; **, $P < 0.01$. **E**, LNCaP and abl cells were transfected with siRNAs in the absence of androgen. Real-time RT-PCR and Western blot analyses were carried out 72 hours after siRNA transfection. **, $P < 0.01$. **F**, ChIP assays were carried out by using an antibody against E2F1 in untreated LNCaP and abl cells. **, $P < 0.01$. **G**, LNCaP and abl cells were transfected with siRNAs in the absence of androgen, and real-time RT-PCR was carried out. *, $P < 0.05$; **, $P < 0.01$.

mRNA expression was greater in abl versus LNCaP in the absence of DHT, showing that *CCNA2* is a direct E2F1 target gene in abl but not LNCaP cells. Thus, FoxA1, through a direct transcriptional regulation of *E2F1*, indirectly upregulates (induces) *CCNA2* expression in abl cells.

Recruitment of MYBL2 and CREB1 leads to FoxA1 binding and CRPC-specific target gene expression through histone acetylation-facilitated nucleosome disruption

We next investigated the mechanisms responsible for abl-specific FoxA1 binding resulting in abl-specific FoxA1 target gene expression. Given that previous studies have reported that cooperation among transcription factors can result in altered chromatin binding activity (7, 11, 16), we hypothesized that this may be the case for FoxA1 binding in abl but not in

LNCaP cells. Thus, to examine whether other transcription factors may affect FoxA1 binding, we conducted a *de novo* transcription factor motif search within the abl- and LNCaP-specific FoxA1 binding regions. As expected, Forkhead motifs were significantly enriched within both abl- and LNCaP-specific FoxA1 binding sites compared with the whole genome background (abl HyperGeometric P values (36): $5.7E-27$ for abl and $1.0E-27$ for LNCaP). Interestingly, MYB and CREB motifs were significantly enriched within abl-specific FoxA1 binding regions compared with LNCaP-specific FoxA1 binding regions (HyperGeometric P values: $5.9E-20$ for MYB and $1.1E-15$ for CREB; Fig. 6A), indicating that transcription factors recognizing MYB and CREB motifs may play a "cooperative role" in FoxA1 binding and abl-specific FoxA1 target gene regulation. As ubiquitous expression of MYBL2 (within the MYB family) and CREB1 (within the CREB family) has been reported




Figure 6. MYBL2 and CREB1 cooperate with FoxA1 to regulate *CCNE2* and *E2F1*. **A**, *de novo* motif search revealed that MYB and CREB motifs are enriched within abl-specific FoxA1 binding sites. **B**, Western blot analyses were carried out on untreated LNCaP and abl cell lysates by using the indicated antibodies. **C**, ChIP assays were carried out by using antibodies against MYBL2 or CREB1 in LNCaP and abl cells in the absence of androgen. *, $P < 0.05$; **, $P < 0.01$. **D–F**, LNCaP and abl cells were transfected with siRNAs in the absence of androgen and ChIP assays were carried out by using an anti-FoxA1 antibody (**D**), or antibodies against MYBL2 or CREB1 (**E**), or an anti-ACh3 antibody (**F**). *, $P < 0.05$; **, $P < 0.01$ as compared with LNCaP or abl siControl. **G**, LNCaP and abl cells were transfected with siRNAs in the absence of androgen and FAIRE-qPCR experiments were carried out. *, $P < 0.05$; **, $P < 0.01$ as compared with LNCaP or abl siControl. **H**, LNCaP and abl cells were transfected with siRNAs in the absence of androgen, and real-time RT-PCR was carried out by using gene-specific primers. *, $P < 0.05$; **, $P < 0.01$. **I**, abl cells were transfected with siRNAs in the absence of androgen, and the cell proliferation was measured, using the WST-1 assay. **, $P < 0.01$.

(37, 38), we examined expression and chromatin binding of MYBL2 and CREB1 in abl and LNCaP cells. MYBL2 and CREB1 protein levels were higher in abl versus LNCaP cells (Fig. 6B), and ChIP assays showed greater recruitment of MYBL2 and CREB1 to regulatory regions of the abl-specific FoxA1 target genes *CCNE2*, *E2F1*, *UBE2C*, and *CDK1* (Fig. 6C; Supplementary Fig. S5A) in abl versus LNCaP cells.

We next examined whether MYBL2 and CREB1 affect FoxA1 binding and p-Pol II loading at regulatory regions of abl-specific FoxA1 target genes. LNCaP and abl cells were transfected with siRNAs targeting MYBL2 or CREB1, and Western blot analyses and ChIP assays were carried out by using an anti-FoxA1 antibody. No effect of MYBL2 or CREB1 silencing on FoxA1 protein expression levels was observed; however, silencing of either transcription factors decreased FoxA1 binding at the enhancers and promoters of *CCNE2* and *E2F1*, as well as the *UBE2C* enhancers and the *CDK1* enhancer in abl but not LNCaP cells (Fig. 6D; Supplementary Fig. S5B

and C). In addition to decreased FoxA1 binding, ChIP analyses further showed that MYBL2 and CREB1 silencing reduced p-Pol II levels on most enhancers and promoters of abl-specific FoxA1 target genes, similar to the effect of FoxA1 silencing itself (Supplementary Fig. S5D). Collectively, these results show that MYBL2 and CREB1 significantly alter FoxA1 binding and p-Pol II loading on chromatin.

To reveal the hierarchical relationship among MYBL2, CREB1 and FoxA1 binding, we silenced FoxA1 and examined MYBL2 and CREB1 binding on chromatin. ChIP analyses showed that FoxA1 silencing decreased MYBL2 and CREB1 recruitment to most regulatory sites of *CCNE2*, *E2F1*, *UBE2C*, and *CDK1* in abl but not LNCaP cells (Fig. 6E; Supplementary Fig. S6A). Serial ChIP (re-ChIP) analyses of FoxA1/MYBL2 and FoxA1/CREB1 further showed a stronger FoxA1-MYBL2 or -CREB1 interaction on abl-specific FoxA1 target gene loci in abl versus LNCaP cells (Supplementary Fig. S6B). These data suggest that FoxA1-MYBL2 or -CREB1 codependently bind to

the same regulatory elements of abl-specific FoxA1 target genes.

To further investigate the molecular mechanisms for the codependent binding of FoxA1-MYBL2 and -CREB1 on chromatin, we examined the effect of their silencing on histone acetylation. Altered chromatin structure can greatly influence transcription factor access to chromatin, and histone acetylation has been shown to have profound effects on chromatin architecture leading to a decrease in internucleosomal interaction (33, 39). LNCaP and abl cells were transfected with siRNAs targeting FoxA1, MYBL2, or CREB1, and ChIP assays were carried out by using an antibody against acetylated histone H3 (AcH3). Silencing of FoxA1, MYBL2, or CREB1 reduced AcH3 levels at regulatory sites of abl-specific FoxA1 target genes in abl but not LNCaP cells, and the level of AcH3 reduction was similar to CBP [a potent histone acetyltransferase (HAT); positive control] silencing (Fig. 6F; Supplementary Fig. S6C), suggesting that these 3 transcription factors, possibly via recruitment of CBP and other HATs, can significantly induce histone acetylation, leading to nucleosome disruption at FoxA1 target gene regulatory regions specifically in abl cells (Figs. 4D, 5D, and 6G; Supplementary Figs. S4B and S6D).

Finally, we examined the effect of MYBL2 and CREB1 silencing on abl-specific FoxA1 target gene expression. Expression of *CCNE2*, *E2F1*, *CCNA2*, *UBE2C*, and *CDK1* after MYBL2 or CREB1 silencing in LNCaP and abl cells were assessed by real-time RT-PCR. Although silencing of MYBL2 and/or CREB1 decreased *CCNE2*, *E2F1*, *CCNA2*, *UBE2C*, and *CDK1* mRNA levels in LNCaP cells to some extent, in abl cells, knocking down of these 2 transcription factors greatly reduced mRNA expression of these abl-specific FoxA1 target genes (Fig. 6H; Supplementary Fig. S6E). As anticipated, slower growth of siMYBL2- or siCREB1-transfected CRPC cells was

observed versus siControl (Fig. 6I; Supplementary Fig. S6F), presumably due to decreased expression of the 5 essential cell-cycle genes, indicating that MYBL2 and CREB1 are critical for abl-specific FoxA1 target gene expression and CRPC cell growth.

Discussion

In this study, we found that in addition to its known role as an AR collaborator in regulating CRPC-specific AR target G_2 -M gene transcription and thus a driver of G_2 -M cell-cycle progression (ref. 11; Fig. 1; Supplementary Figs. S4-S6), FoxA1 directs CRPC G_1 -S cell-cycle progression through direct regulation of *CCNE2*, and indirect regulation of *CCNA2* via *E2F1* (Figs. 2, 4, and 5). These findings establish a previously undescribed yet essential role for FoxA1 as a master cell-cycle regulator, required for G_1 -S and G_2 -M progression in CRPC, whereas AR is mainly involved in promoting cell-cycle G_2 -M but not G_1 -S transition in CRPC (Fig. 7; refs. 11, 40).

The findings that an abl-specific FoxA1 cistrome promotes G_1 -S and G_2 -M cell-cycle progression, through transcriptional regulation of non-AR target genes (*CCNE2* and *E2F1*; Figs. 4 and 5) and AR target genes (*UBE2C* and *CDK1*; Supplementary Fig. S4; ref. 11), raise the question: what are the mechanisms controlling differential FoxA1 binding in abl and LNCaP cells? Although it has been established that H3K4me1 and H3K4 dimethylation (H3K4me2) levels determine differential FoxA1 binding in different cell types (refs. 6, 11; Figs. 4C and 5C), whether differential expression and binding of other transcription factors affect FoxA1 binding is unknown. By employing an integrated computational and experimental approach, we identified transcription factors MYBL2 and CREB1 as potential regulators of abl-specific FoxA1 binding. Furthermore, by combining ChIP, siRNA-ChIP, and re-ChIP analyses,




Figure 7. A model for cell-cycle regulation in CRPC. FoxA1 is a master cell-cycle regulator in CRPC. FoxA1, together with MYBL2 and CREB1, regulates cell-cycle G_1 -S progression through transcriptional regulation of *CCNE2* and *E2F1*. FoxA1 also collaborates with AR to regulate cell-cycle G_2 -M transition through direct regulation of *UBE2C* and *CDK1*.

we showed codependent FoxA1/MYBL2 and FoxA1/CREB1 binding to the same DNA fragments in regulatory regions of abl-specific FoxA1 target genes *CCNE2*, *E2F1*, *UBE2C*, and *CDK1* (Fig. 6; Supplementary Figs. S5 and S6). This codependent chromatin binding is determined by the nonredundant function of FoxA1, MYBL2, and CREB1 in inducing histone H3 acetylation and thus facilitating nucleosome depletion at abl-specific FoxA1 binding regions (Fig. 6F and G; Supplementary Fig. S6C and D). Our findings are consistent with previous *in vitro* findings that activator-dependent histone H3 acetylation leads to an open chromatin structure through disruption of both inter- and intrafiber internucleosome interactions (41, 42). Thus, although histone modifications and nucleosome depletion are not required for FoxA1 to open *in vitro* reconstituted condensed chromatin (4), our findings suggest that *in vivo* FoxA1 binding requires both active histone H3K4 methylation and other collaborating transcription factors capable of inducing histone acetylation and/or nucleosome disruption.

Interestingly, the expression of MYBL2 and CREB1 is not only increased in abl compared with LNCaP (Fig. 6B), but, more importantly, in clinical cases of CRPC versus ADPC

(43, 44). By contrast, strong FoxA1 protein expression persists in early and late phases of prostate cancer (Fig. 1A; refs. 45, 46), suggesting that increased expression and binding of MYBL2 and CREB1 during prostate cancer progression alter FoxA1 genomic binding, leading to CRPC-specific upregulation of critical G₁-S and G₂-M cell-cycle genes by FoxA1 (Fig. 7). Therefore, MYBL2 and CREB1 may serve as new therapeutic targets for CRPC.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Grant Support

This work was supported by NIH grants R00 CA126160 (to Q. Wang and W. Li) and U54 CA113001 (to Q. Wang, K.P. Nephew, and T.H.M. Huang), DOD grant PC094421 (to W. Li), and the Ohio State University Comprehensive Cancer Center (to Q. Wang and T.H.M. Huang).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received June 6, 2011; revised August 4, 2011; accepted August 9, 2011; published OnlineFirst September 7, 2011.

References

- Friedman JR, Kaestner KH. The Foxa family of transcription factors in development and metabolism. *Cell Mol Life Sci* 2006;63:2317–28.
- Gualdi R, Bossard P, Zheng M, Hamada Y, Coleman JR, Zaret KS. Hepatic specification of the gut endoderm in vitro: cell signaling and transcriptional control. *Genes Dev* 1996;10:1670–82.
- Cirillo LA, McPherson CE, Bossard P, Stevens K, Cherian S, Shim EY, et al. Binding of the winged-helix transcription factor HNF3 to a linker histone site on the nucleosome. *EMBO J* 1998;17:244–54.
- Cirillo LA, Lin FR, Cuesta I, Friedman D, Jarnik M, Zaret KS. Opening of compacted chromatin by early developmental transcription factors HNF3 (FoxA) and GATA-4. *Mol Cell* 2002;9:279–89.
- Sekiya T, Muthurajan UM, Luger K, Tulin AV, Zaret KS. Nucleosome-binding affinity as a primary determinant of the nuclear mobility of the pioneer transcription factor FoxA. *Genes Dev* 2009;23:804–9.
- Lupien M, Eeckhoutte J, Meyer CA, Wang Q, Zhang Y, Li W, et al. FoxA1 translates epigenetic signatures into enhancer-driven lineage-specific transcription. *Cell* 2008;132:958–70.
- Hurtado A, Holmes KA, Ross-Innes CS, Schmidt D, Carroll JS. FOXA1 is a key determinant of estrogen receptor function and endocrine response. *Nat Genet* 2011;43:27–33.
- Gao N, Zhang J, Rao MA, Case TC, Mirosevich J, Wang Y, et al. The role of hepatocyte nuclear factor-3 alpha (Forkhead Box A1) and androgen receptor in transcriptional regulation of prostatic genes. *Mol Endocrinol* 2003;17:1484–507.
- Knudsen KE, Penning TM. Partners in crime: deregulation of AR activity and androgen synthesis in prostate cancer. *Trends Endocrinol Metab* 2010;21:315–24.
- Balk SP, Knudsen KE. AR, the cell cycle, and prostate cancer. *Nucl Recept Signal* 2008;6:e001.
- Wang Q, Li W, Zhang Y, Yuan X, Xu K, Yu J, et al. Androgen receptor regulates a distinct transcription program in androgen-independent prostate cancer. *Cell* 2009;138:245–56.
- Culig Z, Hoffmann J, Erdel M, Eder IE, Hobisch A, Hittmair A, et al. Switch from antagonist to agonist of the androgen receptor bicalutamide is associated with prostate tumour progression in a new model system. *Br J Cancer* 1999;81:242–51.
- Sramkoski RM, Pretlow TG 2nd, Giaconia JM, et al. A new human prostate carcinoma cell line, 22Rv1. *In Vitro Cell Dev Biol Anim* 1999;35:403–9.
- Chen Z, Zhang C, Wu D, Chen H, Rorick A, Zhang X, et al. Phospho-MED1-enhanced UBE2C locus looping drives castration-resistant prostate cancer growth. *EMBO J* 2011;30:2405–19.
- Wang H, Zhang C, Rorick A, Wu D, Chiu M, Thomas-Ahner J, et al. CCI-779 inhibits cell-cycle G2-M progression and invasion of castration-resistant prostate cancer via attenuation of UBE2C transcription and mRNA stability. *Cancer Res* 2011;71:4866–76.
- Wang Q, Li W, Liu XS, Carroll JS, Jänne OA, Keeton EK, et al. A hierarchical network of transcription factors governs androgen receptor-dependent prostate cancer growth. *Mol Cell* 2007;27:380–92.
- LaTulippe E, Satagopan J, Smith A, Scher H, Scardino P, Reuter V, et al. Comprehensive gene expression analysis of prostate cancer reveals distinct transcriptional programs associated with metastatic disease. *Cancer Res* 2002;62:4499–506.
- Yu YP, Landsittel D, Jing L, Nelson J, Ren B, Liu L, et al. Gene expression alterations in prostate cancer predicting tumor aggression and preceding development of malignancy. *J Clin Oncol* 2004;22:2790–9.
- Varambally S, Yu J, Laxman B, Rhodes DR, Mehra R, Tomlins SA, et al. Integrative genomic and proteomic analysis of prostate cancer reveals signatures of metastatic progression. *Cancer Cell* 2005;8:393–406.
- Rhodes DR, Yu J, Shanker K, Deshpande N, Varambally R, Ghosh D, et al. ONCOMINE: a cancer microarray database and integrated data-mining platform. *Neoplasia* 2004;6:1–6.
- Eeckhoutte J, Lupien M, Meyer CA, Verzi MP, Shivdasani RA, Liu XS, et al. Cell-type selective chromatin remodeling defines the active subset of FOXA1-bound enhancers. *Genome Res* 2009;19:372–80.
- Giresi PG, Kim J, McDaniell RM, Iyer VR, Lieb JD. FAIRE (Formaldehyde-Assisted Isolation of Regulatory Elements) isolates active regulatory elements from human chromatin. *Genome Res* 2007;17:877–85.
- Liu XS, Brutlag DL, Liu JS. An algorithm for finding protein-DNA binding sites with applications to chromatin-immunoprecipitation microarray experiments. *Nat Biotechnol* 2002;20:835–9.
- Mahony S, Benos PV. STAMP: a web tool for exploring DNA-binding motif similarities. *Nucleic Acids Res* 2007;35:W253–8.
- Sun T, Wang Q, Balk S, Brown M, Lee GS, Kantoff P. The role of microRNA-221 and microRNA-222 in androgen-independent prostate cancer cell lines. *Cancer Res* 2009;69:3356–63.

26. Sobel RE, Sadar MD. Cell lines used in prostate cancer research: a compendium of old and new lines—part 1. *J Urol* 2005;173:342–59.
27. Xu Y, Chen SY, Ross KN, Balk SP. Androgens induce prostate cancer cell proliferation through mammalian target of rapamycin activation and post-transcriptional increases in cyclin D proteins. *Cancer Res* 2006;66:7783–92.
28. Gudas JM, Payton M, Thukral S, Chen E, Bass M, Robinson MO, et al. Cyclin E2, a novel G1 cyclin that binds Cdk2 and is aberrantly expressed in human cancers. *Mol Cell Biol* 1999;19:612–22.
29. Yam CH, Fung TK, Poon RY. Cyclin A in cell cycle control and cancer. *Cell Mol Life Sci* 2002;59:1317–26.
30. Knudsen KE, Arden KC, Cavenee WK. Multiple G1 regulatory elements control the androgen-dependent proliferation of prostatic carcinoma cells. *J Biol Chem* 1998;273:20213–22.
31. Johnson WE, Li W, Meyer CA, Gottardo R, Carroll JS, Brown M, et al. Model-based analysis of tiling-arrays for ChIP-chip. *Proc Natl Acad Sci U S A* 2006;103:12457–62.
32. Heintzman ND, Stuart RK, Hon G, Fu Y, Ching CW, Hawkins RD, et al. Distinct and predictive chromatin signatures of transcriptional promoters and enhancers in the human genome. *Nat Genet* 2007;39:311–8.
33. Cairns BR. The logic of chromatin architecture and remodelling at promoters. *Nature* 2009;461:193–8.
34. Schulze A, Zerfass K, Spitkovsky D, Middendorp S, Bergès J, Helin K, et al. Cell cycle regulation of the cyclin A gene promoter is mediated by a variant E2F site. *Proc Natl Acad Sci U S A* 1995;92:11264–8.
35. Zhu W, Giangrande PH, Nevins JR. E2Fs link the control of G1/S and G2/M transcription. *EMBO J* 2004;23:4615–26.
36. Linhart C, Halperin Y, Shamir R. Transcription factor and microRNA motif discovery: the Amadeus platform and a compendium of meta-zoan target sets. *Genome Res* 2008;18:1180–9.
37. Sala A. B-MYB, a transcription factor implicated in regulating cell cycle, apoptosis and cancer. *Eur J Cancer* 2005;41:2479–84.
38. Mayr B, Montminy M. Transcriptional regulation by the phosphorylation-dependent factor CREB. *Nat Rev Mol Cell Biol* 2001;2:599–609.
39. Kouzarides T. Chromatin modifications and their function. *Cell* 2007;128:693–705.
40. Sharma A, Yeow WS, Ertel A, Coleman I, Clegg N, Thangavel C, et al. The retinoblastoma tumor suppressor controls androgen signaling and human prostate cancer progression. *J Clin Invest* 2010;120:4478–92.
41. Szerlong HJ, Prenni JE, Nyborg JK, Hansen JC. Activator-dependent p300 acetylation of chromatin in vitro: enhancement of transcription by disruption of repressive nucleosome-nucleosome interactions. *J Biol Chem* 2010;285:31954–64.
42. Shahbazian MD, Grunstein M. Functions of site-specific histone acetylation and deacetylation. *Annu Rev Biochem* 2007;76:75–100.
43. Kumar AP, Bhaskaran S, Ganapathy M, Crosby K, Davis MD, Kochunov P, et al. Akt/cAMP-responsive element binding protein/cyclin D1 network: a novel target for prostate cancer inhibition in transgenic adenocarcinoma of mouse prostate model mediated by Nexrutine, a *Phellodendron amurense* bark extract. *Clin Cancer Res* 2007;13:2784–94.
44. Wu D, Zhou HE, Huang WC, Iqbal S, Habib FK, Sartor O, et al. cAMP-responsive element-binding protein regulates vascular endothelial growth factor expression: implication in human prostate cancer bone metastasis. *Oncogene* 2007;26:5070–7.
45. Mirosevich J, Gao N, Gupta A, Shappell SB, Jove R, Matusik RJ. Expression and role of Foxa proteins in prostate cancer. *Prostate* 2006;66:1013–28.
46. Jain RK, Mehta RJ, Nakshatri H, Idrees MT, Badve SS. High-level expression of forkhead-box protein A1 in metastatic prostate cancer. *Histopathology* 2011;58:766–72.